

List of Figures

1.1	A few of the graphical possibilities offered by R	6	2
1.2	The RCommander graphical interface	9	3
1.3	Entering data with the RCommander graphical interface	10	4
1.4	Basic statistics with RCommander	11	5
1.5	Manipulating a data set with RCommander	12	6
1.6	Mean comparison test with RCommander	15	7
1.7	Independence test with RCommander	16	8
1.8	Least squares plane	18	9
3.1	The script window and the command console	41	10
3.2	Characteristics of a complex number	47	11
3.3	Illustration of an array	53	12
7.1	Effect of argument <code>m</code> of function <code>par()</code> . Numbers have been added to gain a better understanding of where future plots will be drawn	13	14
		154	15
7.2	Potential of the function <code>layout()</code>	155	16
7.3	The function <code>layout()</code> and its arguments <code>widths</code> and <code>heights</code>	156	17
7.4	The function <code>plot()</code>	157	18
7.5	The function <code>points()</code>	158	19
7.6	The functions <code>segments()</code> and <code>lines()</code>	159	20
7.7	The function <code>abline()</code>	159	21
7.8	The function <code>arrows()</code>	161	22
7.9	The function <code>curve()</code>	162	23
7.10	The function <code>box()</code>	163	24
7.11	The argument <code>col</code> of function <code>plot()</code>	164	25
7.12	The argument <code>alpha</code> of function <code>rgb()</code>	165	26
7.13	An example using function <code>rainbow()</code>	166	27
7.14	The function <code>display.brewer.all()</code>	167	28
7.15	The function <code>image()</code>	168	29
7.16	The function <code>image()</code> with a coherent display of the data	168	30

7.17	The function <code>text()</code>	169	31
7.18	The function <code>mtext()</code>	170	32
7.19	The function <code>title()</code>	171	33
7.20	Plot title on several lines	172	34
7.21	The function <code>axis()</code>	173	35
7.22	The function <code>legend()</code> with squares	174	36
7.23	The function <code>legend()</code> with line segments	174	37
7.24	Figure illustrating the fine management of graphical parameters ...	178	38
7.25	Managing the colours of a plot	179	39
7.26	Example of use of the arguments <code>adj</code> and <code>srt</code>	181	40
7.27	Using different fonts on a plot	182	41
7.28	Labels on a plot	184	42
7.29	The arguments <code>lend</code> and <code>ljoin</code>	185	43
7.30	The argument <code>pch</code>	186	44
7.31	The arguments <code>lty</code> and <code>lwd</code>	186	45
8.1	Result of the call of the function <code>mydisplay.reg1()</code>	211	46
8.2	Emacs and GDB	263	47
8.3	DDD and GDB	265	48
9.1	Illustration of storage of values in memory. Each little box contains a binary number (0 or 1). Each green number gives the decimal representation of the number in binary form in the block above. Each red number gives the address (expressed here in decimal notation) of the 8-box block above. Note that the same memory addresses could have been written in hexadecimal notation ($b = 16$), giving 3C, 3D, 3E and 3F	294	49 50 51 52 53 54 55
9.2	Illustration of R storage in memory of a (signed) integer. Each little box contains a binary digit (0 or 1). The green number gives the decimal notation of the integer expressed in binary notation in the four blocks above. The red number gives the address (expressed here in decimal base) of the first 8-box memory block above. Note that here, a number is stored over 32 boxes and not over 8 as in Fig. 9.1. Furthermore, the first box is used to specify the sign of the number, negative here	295	56 57 58 59 60 61 62 63
10.1	Modified sinc function	329	64
11.1	Algorithm to determine the type of a variable	340	65
11.2	Cross chart for a qualitative variable	358	66
11.3	Dot chart for a qualitative variable	358	67
11.4	Bar chart for a qualitative variable	359	68
11.5	Pareto chart for a qualitative variable	360	69
11.6	Stacked bar chart for a qualitative variable	361	70
11.7	Bar chart with cumulative frequencies line for an ordinal variable	363	71 72

11.8	Bar chart for a discrete quantitative variable	364	73
11.9	Empirical distribution function for a discrete quantitative variable..	364	74
11.10	Boxplot and explanations	366	75
11.11	Plot of the empirical distribution function for a continuous quantitative variable.	367	76 77
11.12	Density histogram with identical or different class ranges	369	78
11.13	Frequency polygon	370	79
11.14	Cumulative frequency polygon	371	80
11.15	Bar plot for two qualitative variables	372	81
11.16	Mosaic plot for two qualitative variables	372	82
11.17	Cohen–Friendly association plot for two qualitative variables	373	83
11.18	table. cont plot for two qualitative variables	374	84
11.19	Plot of two quantitative variables	375	85
11.20	Box plots of a quantitative variable as a function of the levels of a qualitative variable	376	86 87
11.21	stripchart plot for a quantitative and a qualitative variable.	376	88
12.1	Plot approximating the density of X	388	89
12.2	Convergence in distribution in action on an example with simulated data.	394	90 91
14.1	Scatter plot of child weight against mother weight.	457	92
14.2	Least squares regression line on a scatter plot.	458	93
14.3	Visualization of confidence and prediction intervals	463	94
14.4	Graphical inspection of normality of residuals	464	95
14.5	Plot of residuals as a function of predicted values	465	96
14.6	Scatter plot of all pairs of variables	470	97
14.7	Effect of age on BWT in a model without interaction	479	98
14.8	Effect of age on BWT in a model with interaction	480	99
14.9	Selecting variables with the BIC	484	100
14.10	Checking the assumptions of homoscedasticity (<i>left</i>) and normality (<i>right</i>)	490	101 102
14.11	Residuals as a function of explanatory variables	491	103
14.12	Visualizing outliers: studentized residuals against fitted values	492	104
14.13	Visualizing influential observations: Cook’s distance	494	105
15.1	Box plot of scarring times for each treatment	506	106
15.2	Analysing the residuals in single-factor ANOVA	509	107
15.3	Exploration of interaction in two-way ANOVA	516	108
15.4	Residual analysis in two-way ANOVA	520	109

UNCORRECTED PROOF